



Whitsunday Water Quality Monitoring Blueprint for Tourism Operators: Quarterly update - October 2020

Jordan Iles and Nathan Waltham

Report No. 20/35

October 2020



Whitsunday Water Quality Monitoring Blueprint for Tourism Operators: Quarterly update - October 2020

A Report for Reef Catchments (Mackay Whitsunday Isaac) Limited

Report No. 20/35

October 2020

Prepared by Jordan Iles and Nathan Waltham

[Centre for Tropical Water & Aquatic Ecosystem Research](#)

[\(TropWATER\)](#)

James Cook University

Townsville

Phone : (07) 4781 4262

Email: TropWATER@jcu.edu.au

Web: www.jcu.edu.au/tropwater/

Information should be cited as:

Iles, JA & Waltham, NJ 2020, 'Whitsunday Water Quality Monitoring Blueprint for Tourism Operators: Quarterly update - October 2020', Centre for Tropical Water & Aquatic Ecosystem Research (TropWATER) Publication, James Cook University, Townsville, 19 pp.

For further information contact:

Dr Nathan Waltham
Centre for Tropical Water & Aquatic Ecosystem Research (TropWATER)
James Cook University
nathan.waltham@jcu.edu.au

This publication has been compiled by the Centre for Tropical Water & Aquatic Ecosystem Research (TropWATER), James Cook University.

© James Cook University, 2020.

Except as permitted by the *Copyright Act 1968*, no part of the work may in any form or by any electronic, mechanical, photocopying, recording, or any other means be reproduced, stored in a retrieval system or be broadcast or transmitted without the prior written permission of TropWATER. The information contained herein is subject to change without notice. The copyright owner shall not be liable for technical or other errors or omissions contained herein. The reader/user accepts all risks and responsibility for losses, damages, costs and other consequences resulting directly or indirectly from using this information.

Enquiries about reproduction, including downloading or printing the web version, should be directed to nathan.waltham@jcu.edu.au

Acknowledgments:

The Whitsunday Water Quality Monitoring Blueprint for Tourism Operators is funded by the partnership between the Australian Government's Reef Trust and the Great Barrier Reef Foundation. Cover photo provided by Reef Catchments.



Great Barrier
Reef Foundation

TABLE OF CONTENTS

TABLE OF FIGURES.....	4
1 INTRODUCTION.....	5
2 RESULTS.....	6
2.1 Logger data.....	6
2.1.1 Water temperature.....	6
2.1.2 Water depth and wave height.....	7
2.1.3 Photosynthetically active radiation (PAR).....	9
2.1.4 Turbidity.....	11
2.2 Water samples.....	12
2.2.1 Field observations.....	12
2.2.2 Physico-chemical parameters.....	13
2.2.3 Nutrients.....	15
2.2.4 Chlorophyll <i>a</i>	18

TABLE OF FIGURES

Figure 1. Water temperature measured at Cairn Beach and Tongue Bay 2020.....	6
Figure 2. Water depth measured at Cairn Beach and Tongue Bay 2020.....	7
Figure 3. Root mean squared (RMS) water depth measured at Cairn Beach and Tongue Bay 2020.....	8
Figure 4. Photosynthetically active radiation (PAR) measured at 10 min intervals at Cairn Beach and Tongue Bay 2020.....	9
Figure 5. Daily photosynthetically active radiation at Cairn Beach and Tongue Bay 2020.....	10
Figure 6. Turbidity measured at Cairn Beach and Tongue Bay 2020. Note different y-axis scale. Data has been removed from the timeseries for periods where the sensor fouled.....	11
Figure 7. Secchi depth measured at Cairn Beach and Tongue Bay 2020.....	12
Figure 8. Electrical conductivity (Cond), total suspended solids (TSS), and pH of water samples collected from Cairn Beach WH1 (green) and Tongue Bay WH2 (blue) 2020.....	14
Figure 9. Total nitrogen (TN), total dissolved nitrogen (TDN), particulate nitrogen (PN), and nitrate-nitrite (NO _x) concentrations measured in water samples collected from Cairn Beach WH1 (green) and Tongue Bay WH2 (blue) 2020.....	16
Figure 10. Total phosphorus (TP), total dissolved phosphorus (TDP), and particulate phosphorus (PP) concentrations measured in water samples collected from Cairn Beach WH1 (green) and Tongue Bay WH2 (blue) 2020.....	17
Figure 11. Chlorophyll <i>a</i> and phaeophytin <i>a</i> concentrations measured in water samples collected from Cairn Beach WH1 (green) and Tongue Bay WH2 (blue) 2020.....	19

1 INTRODUCTION

This report gives a quarterly update on data collected by the Whitsunday Water Quality Monitoring Blueprint for Tourism Operators in the third quarter of 2020. The water sampling and logger maintenance schedule for this period is outlined in Table 1.

For project background and methodology please refer to the annual report: Iles, JA & Waltham, NJ 2020, 'Whitsunday Water Quality Monitoring Blueprint for Tourism Operators: Annual report 2019-2020', Centre for Tropical Water & Aquatic Ecosystem Research (TropWATER) Publication, James Cook University, Townsville, 29 pp.

Table 1. Monitoring events for the third quarter of 2020. Events covered in previous reports are shown in grey.

Event	Date scheduled	Cairn Beach (WH1)	Tongue Bay (WH2)
a	04/02/2020	Ocean Rafting / TropWATER	Ocean Rafting / TropWATER
b	04/03/2020	True Blue	Red Cat
c	11/05/2020	Ocean Rafting	Ocean Rafting
d [#]	26/05/2020	Thunder Cat	Red Cat
e	23/06/2020	Ocean Rafting	Ocean Rafting
f	08/08/2020	Ocean Rafting	Ocean Rafting
g	17/09/2020	Ocean Rafting / TropWATER	Ocean Rafting / TropWATER

[#] water sampling only

2 RESULTS

The results within this report cover the period 24/06/2020 to 17/09/2020. There were two logger instrument deployments during this period. Deployment 4 covered 24/06/2020 to 08/08/2020 (46 days), and Deployment 5 covered the period to 08/08/2020 to 17/09/2020 (40 days). Results reported in previous reports are shown in grey on figures and tables throughout.

2.1 Logger data

2.1.1 Water temperature

Water temperature during the last two deployment periods ranged from 21.1 to 23.2 °C at Cairn Beach and 20.6 to 23.4 °C at Tongue Bay (Figure 1). Water temperatures continued to cool until August, and have since been warming as we come into spring.




Figure 1. Water temperature measured at Cairn Beach and Tongue Bay 2020

2.1.2 Water depth and wave height

The average water depths of the logger at Cairn Beach were 11.0 and 10.0 m for Deployments 4 and 5 respectively. Average water depths of the logger at Tongue Bay were 8.7 and 7.4 m for Deployments 4 and 5 respectively (Figure 2). It is apparent from the depth that that the position of the loggers changed between deployments, with both the Cairn beach and Tongue Bay loggers sitting in slightly shallower water during the fifth deployment. The root means square (RMS) depth is shown in Figure 3. There was very little wave activity at the Cairn Beach site with RMS depth generally less than 0.005 m throughout the deployment periods (average = 0.003 m). Tongue Bay was more exposed to wave energy with a mean RMS depth of 0.020 m.




Figure 2. Water depth measured at Cairn Beach and Tongue Bay 2020




Figure 3. Root mean squared (RMS) water depth measured at Cairn Beach and Tongue Bay 2020

2.1.3 Photosynthetically active radiation (PAR)

Photosynthetically active radiation (PAR) measured at depth followed a typical diel cycle with higher values during the day, decreasing to zero overnight (Figure 4). Daily photosynthetically active radiation is shown in Figure 5. Mean daily PAR values were 1.56 and 1.0 mol m⁻² d⁻¹ at Cairn Beach, and 0.44 and 1.04 mol m⁻² d⁻¹ at Tongue Bay for Deployments 4 and 5 respectively.




Figure 4. Photosynthetically active radiation (PAR) measured at 10 min intervals at Cairn Beach and Tongue Bay 2020




Figure 5. Daily photosynthetically active radiation at Cairn Beach and Tongue Bay 2020

2.1.4 Turbidity

Turbidity ranged from 0.05 to 26.5 NTUe at the Cairn Beach monitoring site, with an average value of 1.04 NTUe over the deployment period (Figure 6). Turbidity ranged from 0 to 33.8 NTUe at the Tongue Bay monitoring site, with an average value of 6.38 NTUe over the deployment period. Both the Cairn Beach and Tongue Bay instruments had issues with sensor fouling during the deployment, with biofouling present on the sensor when inspected post-deployment. Turbidity data for Tongue Bay was deemed unreliable due to sensor fouling after 7th June (day 14) for deployment 4, and after 28th August (day 20) for deployment 5 and removed during the data QC process.




Figure 6. Turbidity measured at Cairn Beach and Tongue Bay 2020. Note different y-axis scale. Data has been removed from the timeseries for periods where the sensor fouled.

2.2 Water samples

2.2.1 Field observations

The tourism operators recorded observations of conditions on the water at the time of each sampling and maintenance event (Table 2). Secchi depths measured by tourism operators at Cairn Beach and Tongue Bay are shown in **Figure 7**.

Table 2. Water quality measurements and observations recorded from Cairn Beach and Tongue Bay.

Site Name	Site Code	Date	Time	Secchi depth m	Cloud cover %	Wind knots	Sea surface	Surface scum/slick
Cairn Beach	WH1	04/02/2020	13:45	4.5	15	5 - 10	Calm	Nil
		07/03/2020	17:15	4	80	0 - 5	Flat	Nil
		11/05/2020	08:05	3	75	21 - 25	Light surface chop	Nil
		26/05/2020	11:45	6	5 - 10	5 - 10	Calm	Nil
		23/06/2020	08:15	6	0	5 - 10	Calm	Nil
		08/08/2020	09:15	5.1	100	0	0	Nil
		17/09/2020	09:40	3	70	10 - 15	Choppy	Nil
Tongue Bay	WH2	04/02/2020	15:00	6.5	10	5 - 10	Calm	Nil
		09/03/2020	10:30	0.3 [#]	80	5	Glass	Nil
		11/05/2020	10:20	3 [*]	5	25	Chop	Nil
		26/05/2020	12:30	5.5	60	5	Calm	Nil
		23/06/2020	09:30	4.2	0	10 - 15	0 - 0.3	Nil
		08/08/2020	10:15	5	80	5	Slight, Ripple	Nil
		17/09/2020	11:20	2.9	90	10 - 15	Choppy	Nil

[#] Note: Likely erroneous measurement, * Note: Secchi depth estimated




Figure 7. Secchi depth measured at Cairn Beach and Tongue Bay 2020

2.2.2 Physico-chemical parameters

Water samples collected from the monitoring sites during August and September 2020 were measured in the laboratory for electrical conductivity, total suspended solids (TSS), and pH (Table 3, Figure 8). Conductivity ranged from 52.95 to 53.70 mS cm⁻¹ and was within expected range of seawater with limited freshwater inputs. TSS ranged from 1.0 to 2.0 mg L⁻¹. pH ranged from 8.14 to 8.19.

Table 3. Electrical conductivity (Cond, mS cm⁻¹), Total suspended solids (TSS, mg L⁻¹) and pH of surface water samples collected from Cairn Beach and Tongue Bay.

Site name	Site code	Sample Date	Cond mS cm ⁻¹	TSS mg L ⁻¹	pH
Cairn Beach	WH1	4/02/2020	54.02	1.9	8.29
		9/03/2020	52.95	9.3	8.33
		11/05/2020	53.03	2.2	8.35
		26/05/2020	53.08	1.9	8.24
		23/06/2020	53.62	2.4	8.25
		8/08/2020	52.96	1.0	8.19
		17/09/2020	53.63	2.0	8.14
Tongue Bay	WH2	4/02/2020	53.92	2.8	8.31
		8/03/2020	53.14	3.0	8.29
		11/05/2020	53.07	2.0	8.32
		26/05/2020	53.01	1.8	8.27
		23/06/2020	53.64	2.6	8.23
		8/08/2020	52.95	1.1	8.14
		17/09/2020	53.70	1.7	8.15




Figure 8. Electrical conductivity (Cond), total suspended solids (TSS), and pH of water samples collected from Cairn Beach WH1 (green) and Tongue Bay WH2 (blue) 2020.

2.2.3 Nutrients

Results for water samples collected during the two most recent scheduled trips are presented below, with previous sampling events shown in grey for comparison. Nitrogen and phosphorus concentrations are presented in Table 4. Total nitrogen concentrations ranged from 94 to 254 $\mu\text{g N L}^{-1}$ during the most recent sampling events (Figure 9). Total phosphorus concentrations ranged from 6 to 7 $\mu\text{g P L}^{-1}$ (Figure 10). The bulk of nutrients were present in the dissolved fraction with 50 to 93% nitrogen dissolved, and 71 to 83 % phosphorus dissolved. Nitrate-nitrite concentrations ranged 2 to 5 $\mu\text{g N L}^{-1}$.

Table 4. Nutrient concentrations measured in surface water samples collected from Cairn Beach and Tongue Bay. Total nitrogen (TN), total dissolved nitrogen (TDN), particulate nitrogen (PN), nitrate-nitrite (NO_x), total phosphorus (TP), total dissolved phosphorus (TDP), and particulate phosphorus (PP). All concentrations reported in micrograms per litre ($\mu\text{g L}^{-1}$).

Site name	Site code	Sample Date	TN $\mu\text{g L}^{-1}$	TDN $\mu\text{g L}^{-1}$	PN $\mu\text{g L}^{-1}$	NO_x $\mu\text{g L}^{-1}$	TP $\mu\text{g L}^{-1}$	TDP $\mu\text{g L}^{-1}$	PP $\mu\text{g L}^{-1}$
Cairn Beach	WH1	4/02/2020	142	136	6	6	9	8	1
		9/03/2020	227	111	116	7	17	7	10
		11/05/2020	158	130	28	2	7	6	1
		26/05/2020	93	84	9	3	6	5	1
		23/06/2020	85	84	1	3	6	6	0
		8/08/2020	160	114	46	5	7	6	1
		17/09/2020	113	82	31	5	6	3	3
Tongue Bay	WH2	4/02/2020	126	118	8	3	9	8	1
		8/03/2020	130	109	21	3	8	7	1
		11/05/2020	169	155	14	2	8	6	2
		26/05/2020	102	82	20	2	7	5	2
		23/06/2020	109	88	21	5	6	5	1
		8/08/2020	254	126	128	3	7	5	2
		17/09/2020	94	87	7	2	6	5	1




Figure 9. Total nitrogen (TN), total dissolved nitrogen (TDN), particulate nitrogen (PN), and nitrate-nitrite (NOx) concentrations measured in water samples collected from Cairn Beach WH1 (green) and Tongue Bay WH2 (blue) 2020.




Figure 10. Total phosphorus (TP), total dissolved phosphorus (TDP), and particulate phosphorus (PP) concentrations measured in water samples collected from Cairn Beach WH1 (green) and Tongue Bay WH2 (blue) 2020.

2.2.4 Chlorophyll *a*

Chlorophyll *a* values ranged from below detection limit ($<0.2 \mu\text{g L}^{-1}$) to $0.49 \mu\text{g L}^{-1}$ between sites during the most recent sampling events (Table 5, Figure 11). Phaeophytin *a* concentrations were below detection limit ($< 0.2 \mu\text{g L}^{-1}$).

Table 5. Chlorophyll-*a* and Phaeophytin-*a* concentrations measured in surface water samples collected from Cairn Beach and Tongue Bay. All concentrations reported in micrograms per litre ($\mu\text{g L}^{-1}$).

Site name	Site code	Sample Date	Chl- <i>a</i> $\mu\text{g L}^{-1}$	Phaeo- <i>a</i> $\mu\text{g L}^{-1}$
Cairn Beach	WH1	4/02/2020	0.41	<0.2
		9/03/2020	0.38	<0.2
		11/05/2020	0.39	<0.2
		26/05/2020	0.46	0.23
		23/06/2020	0.28	0.25
		8/08/2020	<0.2	<0.2
		17/09/2020	0.45	<0.2
Tongue Bay	WH2	4/02/2020	0.40	0.22
		8/03/2020	0.40	0.21
		11/05/2020	0.44	<0.2
		26/05/2020	0.53	<0.2
		23/06/2020	0.25	<0.2
		8/08/2020	<0.2	<0.2
		17/09/2020	0.49	<0.2




Figure 11. Chlorophyll *a* and phaeophytin *a* concentrations measured in water samples collected from Cairn Beach WH1 (green) and Tongue Bay WH2 (blue) 2020.